$$c = 3.10^8 \,\text{m/s}$$
;  $h = 6.63.10^{-14} \,\text{J} \cdot \text{s}$ ;  $1 \,\text{eV} = 1.60.10^{-19} \,\text{J}$ 

42) The Sun is approximately an ideal blackbody radiator with a surface temperature of 5800 K. (a) Find the wavelength at which its spectral radiancy is maximum and (b) identify the type of electromagnetic wave corresponding to that wavelength. (See Fig. 33-1.) (c) The universe is approximately an ideal blackbody radiator with radiation emitted when atoms first formed. Today the spectral radiancy of that radiation peaks at a wavelength of 1.06 mm (in the microwave region). What is the corresponding temperature of the universe?




Figure 33-1 The electromagnetic spectrum.

- 43) Just after detonation, the fireball in a nuclear blast is approximately an ideal blackbody radiator with a surface temperature of about  $1.0 \cdot 10^7$  K. (a) Find the wavelength at which the thermal radiation is maximum and (b) identify the type of electromagnetic wave corresponding to that wavelength. (See Fig. 33-1.) This radiation is almost immediately absorbed by the surrounding air molecules, which produces another ideal blackbody radiator with a surface temperature of about  $1.0 \cdot 10^5$  K. (c) Find the wavelength at which the thermal radiation is maximum and (d) identify the type of electromagnetic wave corresponding to that wavelength.
- 15) Light strikes a sodium surface, causing photoelectric emission. The stopping potential for the ejected electrons is 5.0 V, and the work function of sodium is 2.2 eV. What is the wavelength of the incident light?
- 61) The function  $\psi(x)=Ae^{ikx}$  can describe a free particle, for which the potential energy is U(x)=0 in Schrödinger's equation. Assume now that  $U(x)=U_0=0$  a constant in that equation. Show that  $\psi(x)$  is a solution of Schrödinger's equation

$$\frac{d^2\psi}{dx^2} + \frac{8\pi^2 m}{h^2} [E - U(x)]\psi = 0$$

for a specific value of the angular wave number k of the particle.